

Multivariate Time Series: Part 3

Identification and Estimation of Simultaneous Equations Systems

Gerald P. Dwyer

Clemson University

March 2021

Outline

1 Multivariate Time Series

- Review
- Structural Equations and Identification
- Estimation Methods
- Generalized Method of Moments
- GMM When Parameters Overidentified
- Summary

Nonunique Representation

- Even more obviously than in univariate time series, there are various observationally equivalent representations of time series
- Normalization in the context of a vector moving-average representation

$$\mathbf{y}_t = \mathbf{B}(L) \boldsymbol{\varepsilon}_t$$

- Normalization issue in terms of vector autoregressions (VARs)

$$\mathbf{A}_0 \mathbf{y}_t = \mathbf{A}_1(L) \mathbf{y}_{t-1} + \boldsymbol{\varepsilon}_t$$

- ▶ Recursive dynamic systems, for example,

$$y_{1,t} = a_{1,2}^* y_{2,t} + a_{1,1} y_{1,t-1} + a_{1,2} y_{2,t-1} + \varepsilon_{1,t}$$

$$y_{2,t} = a_{2,1} y_{1,t-1} + a_{2,2} y_{2,t-1} + \varepsilon_{2,t}$$

Systems of behavioral equations

- Relationship of VAR to economic system

$$q_t = \alpha p_t + \beta y_t + \varepsilon_t^d, \quad \alpha < 0, \beta > 0$$

$$q_t = \gamma p_t + \delta w_t + \varepsilon_t^s, \quad \gamma, \delta > 0$$

Structural equations

- Simultaneous system for supply and demand

$$q_t = \alpha p_t + \beta y_t + \varepsilon_t^d, \alpha < 0, \beta > 0$$

$$q_t = \gamma p_t + \delta w_t + \varepsilon_t^s, \gamma, \delta > 0$$

- Common usage in macroeconomics: An equation is *structural* relative to an intervention if it is invariant to that intervention
- Example of intervention
 - ▶ Suppose a sales tax is imposed on purchases and the demand equation is changed to include $(1 + \tau) p$ where τ is the sales tax
 - ▶ Then these equations are structural relative to the intervention of a change in τ

Model, structure and observational equivalence

- Simultaneous system for supply and demand

$$q_t = \alpha p_t + \beta y_t + \varepsilon_t^d, \varepsilon_t^d \sim N(0, \sigma_d^2)$$

$$q_t = \gamma p_t + \delta w_t + \varepsilon_t^s, \varepsilon_t^s \sim N(0, \sigma_s^2)$$

- This is a **model**
- A **structure** based on this model includes the model and a set of parameter values, including the variances of the innovations
- A structure is **identified** if a structure generates a one-to-one mapping from parameter values to the probability distribution of the observables
- Two structures are **observationally equivalent** if they generate the same probability distribution of the observables
- A model is identified if all possible (or at least interesting) structures are identified
- If no structure is identified, then the model is under-identified or not identified

Example of model, structure and observational equivalence

- Model

$$\alpha_0 q_t = \alpha p_t + \beta y_t + \gamma_0 \varepsilon_t^d, \quad \varepsilon_t^d \sim N(0, \sigma_d^2)$$

- A structure is

$$2q_t = 4p_t + 3y_t + 9\varepsilon_t^d, \quad \varepsilon_t^d \sim N(0, 5)$$

- This structure is not identified because the identical set of pdfs of the variables is generated by

$$4q_t = 8p_t + 6y_t + 18\varepsilon_t^d, \quad \varepsilon_t^d \sim N(0, 5)$$

- Identify it by

- ▶ Normalizing on a dependent variable, e.g. $\alpha_0 = 1$ or $\alpha = -1$
 - ★ “For example” because could normalize to some other nonzero value
- ▶ And normalizing the coefficient of the error term γ_0 to $\gamma_0 = 1$ or $\gamma_0 = -1$

Identification of simultaneous-equation system

- In the simultaneous system for supply and demand, we solved for the reduced form

$$q = (\gamma - \alpha)^{-1} \left[\gamma\beta y - \alpha\delta w + \gamma\varepsilon_t^d - \alpha\varepsilon_t^s \right]$$
$$p = (\gamma - \alpha)^{-1} \left[\beta y - \delta w + \varepsilon_t^d - \varepsilon_t^s \right]$$

- ▶ which also can be written

$$\begin{aligned} q &= \pi_{11}y + \pi_{12}w + u_1 \\ p &= \pi_{21}y + \pi_{22}w + u_2 \end{aligned} \tag{1}$$

- Can solve for the parameters in the simultaneous equations uniquely from the reduced form
- Reduced form can be estimated consistently by OLS and is identified
- Therefore the simultaneous equations are identified

Identification of simultaneous-equation system

- Simultaneous system for supply and demand

$$q_t = \alpha p_t + \beta y_t + \beta_2 w_t + \varepsilon_t^d$$

$$q_t = \gamma p_t + \delta w_t + \delta_2 y_t + \varepsilon_t^s$$

- This set of equations is an example of a model that is not identified
 - ▶ You can see this by solving for reduced form and trying to solve back for structural parameters from reduced form (Do yourself)
 - ▶ There are four reduced-form coefficients and six structural coefficients
- In general, how do we identify coefficients?

Restrictions to identify simultaneous-equation system

- Normalizations
 - ▶ Example: coefficients on some variables equal to one
- Use linear identities in model
 - ▶ Examples

$$c + i + g + nx = y$$

$$R = r + \pi$$

$$m = M - P$$

- Exclusion restrictions
 - ▶ Example: coefficient of w_t is zero in demand equation and coefficient of y_t is zero in supply equation

Restrictions to identify simultaneous-equation system

- Restrictions on coefficients
 - ▶ Value of parameters known or restrictions on linear functions of coefficients
 - ▶ Example: sum of coefficients equal to one
- Restrictions on structural variance-covariance matrix
 - ▶ Example: uncorrelated errors across equations
- Nonlinear functions
 - ▶ Example: one variable is a nonlinear function of another

Exclusion restrictions

- Order condition: The number of exogenous variables excluded from an equation is at least as large as the number of jointly dependent variables included in the equation minus one
 - ▶ Counts all jointly dependent variables in the equation
 - ▶ Over-identified if the number of exogenous variables excluded is greater than the number of jointly dependent variables included in the equation minus one
- Rank condition: A restriction on a sub-matrix of the reduced-form coefficients

Illustration of identification

- Shifts in the demand curve identify supply curve
- Shifts in the supply curve identify demand curve
 - ▶ Famous papers by E. J. Working (1927) and T. C. Koopmans (1949)

Application of exclusion restrictions: order condition

- Order condition: The number of exogenous variables excluded from an equation is at least as large as the number of jointly dependent variables included in the equation minus one
- Simultaneous system for supply and demand

$$q_t = \alpha p_t + \beta y_t + \varepsilon_t^d, \quad \alpha < 0, \beta > 0$$

$$q_t = \gamma p_t + \delta w_t + \varepsilon_t^s, \quad \gamma > 0, \delta > 0$$

- Another simultaneous system for supply and demand

$$q_t = \alpha p_t + \beta y_t + \beta_1 z_t + \varepsilon_t^d, \quad \alpha < 0, \beta > 0$$

$$q_t = \gamma p_t + \delta w_t + \varepsilon_t^s, \quad \gamma > 0, \delta > 0$$

- Yet another simultaneous system for supply and demand

$$q_t = \alpha p_t + \varepsilon_t^d, \quad \alpha < 0$$

$$q_t = \gamma p_t + \delta w_t + \varepsilon_t^s, \quad \gamma > 0, \delta > 0$$

Estimation Methods

- There are a variety of ways to estimate structural equations
- What follows is an informal survey of estimation methods with discussion
 - ▶ Indirect least squares (ILS)
 - ▶ Instrumental variables (IV)
 - ▶ Two-stage least squares (2SLS)
 - ▶ Generalized method of moments (GMM)
 - ▶ Limited-information maximum likelihood (LIML)
 - ▶ System methods: Three-stage least squares (3SLS) and Full-information maximum likelihood (FIML)
- Exposition of GMM

Indirect least squares

- We can estimate the parameters by estimating the reduced form by OLS and then inferring the structural parameters
- This is called **indirect least squares**
 - ▶ Estimate reduced form by OLS and solve for structural parameters
 - ▶ With simple equations here, can estimate reduced form consistently by ordinary least squares
 - ▶ Given the estimated reduced form, solve for the structural coefficients
- Works if structural equation is exactly identified
 - ▶ If under-identified, cannot solve uniquely for structural parameters
 - ▶ If over-identified, there is more than one implied value of one or more parameters
 - ★ How choose which one?

OLS and estimation of structural equations

- Why can't we just estimate the structural equations by OLS?
- We know from last lecture that OLS is inconsistent
- People often use OLS to estimate a structural equation to get a “feel” for the correlations in the data
 - ▶ What are the effects of the complicated estimation strategy?
 - ▶ Not a substitute for estimating the parameters in a consistent way!

Instrumental variables

- Instrumental variables here is no different than instrumental variables in other contexts
- Want variables not included in the equation that are correlated with the variable of interest and not with the error term in the equation
- Probably most common estimation strategy used

Instrumental variables and exogenous variables

- Why not use exogenous variables in system as instruments?
- By assumption, uncorrelated with error term
- By assumption, correlated with jointly dependent variables

Two-stage least squares

- Two-stage least squares (2SLS) uses all the exogenous variables in the system of equations as instruments
- By assumption these variables are uncorrelated with the error term in the equation being estimated
- Regress right-hand-side jointly dependent variables on all the exogenous variables
- Use predicted values in structural equation instead of actual values of right-hand-side jointly dependent variables
- Run OLS to estimate coefficients with predicted values
 - ▶ Not to estimate standard errors of coefficients!
- Assumes homoskedastic errors in equations

Generalized method of moments

- Generalized method of moments (GMM) is similar to IV and 2SLS
- Can allow for heteroskedastic errors
- Easier to include covariance restrictions
- Will go through GMM in more detail later in lecture

Limited-information maximum likelihood

- Limited-information maximum likelihood (LIML) maximizes the likelihood for one equation
- Assume likelihood for errors (generally normal) and maximize likelihood
- Same asymptotic distribution as 2SLS if normal distribution assumption is satisfied
- Advantage over 2SLS
 - ▶ 2SLS generally produces different estimates with different normalizations of the equation (Why?)
 - ▶ LIML estimates are unaffected by the normalization of the equation

System methods

- Three-stage least squares (3SLS) and Full-information maximum likelihood (FIML)
- 3SLS – Similar to Seemingly Unrelated Regressions with 2SLS estimates
- FIML – Just maximize likelihood function for system

Single equation versus system

- In system methods, a specification error in one equation is propagated throughout the entire set of equations

Method of Moments

- Generalized method of moments is a generalization of the Method of Moments
- Method of moments
 - ▶ Use moments of data to estimate parameters
 - ▶ Suppose want to estimate n parameters characterizing a distribution
 - ▶ Suppose there exists a well-defined mapping from n parameters θ_i to the first n moments μ_i of the distribution of a variable y
 - ▶ This can be written as

$$\mu_1 = f(\theta_1, \theta_2, \dots, \theta_n)$$

$$\mu_2 = f(\theta_1, \theta_2, \dots, \theta_n)$$

...

$$\mu_n = f(\theta_1, \theta_2, \dots, \theta_n)$$

- ▶ Method of moments estimators substitute sample moments for population moments and estimate parameters by solving the equations

Method of Moments

- Simple examples of Method of Moments
- Want to compute expected value $E y_t$
 - ▶ Use mean for expected value

$$E y_t = \mu_1$$

- Use first and second moment to compute expected value and variance

$$E y_t = \mu_1$$
$$\text{Var}[y_t] = E [y_t - E y_t]^2 = \mu_2 - \mu_1^2$$

or

$$\mu_1 = E y_t$$

$$\mu_2 = \text{Var}[y_t] + \mu_1^2$$

- MOM involves substitution of the sample moments for the population moments in these equations

Method of Moments

- MOM involves substitution of the sample moments for the population moments
- Under fairly general conditions, MOM is a consistent estimator of the parameters of interest

Generalized Method of Moments I

- Can think of moments more generally
- Moments (uncentered) in general are

$$E y_t = \mu_1$$

$$E y_t^2 = \mu_2$$

...

$$E y_t^n = \mu_n$$

- Think of conditional expectations and other functions of the data
- Regression example

$$y_t = \beta x_t + \varepsilon_t$$

- ▶ Impose restriction that

$$E [\varepsilon_t | x_t] = 0$$

- ▶ Limit analysis to linear projections, or normal distributions

Generalized Method of Moments II

- ▶ For linear projections,

$$E[\varepsilon_t | x_t] = \text{Cov}[\varepsilon_t, x_t]$$

- ▶ Given zero means

$$E[\varepsilon_t | x_t] = E \varepsilon_t x_t$$

- ▶ Proceed by imposing restriction in the sample equation

$$y_t = b x_t + e_t$$

$$E e_t x_t = 0$$

which is the same as imposing that the covariance of e_t and x_t is zero

GMM and Ordinary Least Squares I

- The covariance restriction is

$$E[\varepsilon_t x_t] = 0$$

- This implies

$$\begin{aligned} \sum_{t=1}^T e_t x_t &= 0 \\ &= \sum_{t=1}^T (y_t - b x_t) x_t \\ &= \sum_{t=1}^T y_t x_t - \sum_{t=1}^T b x_t x_t \\ &= \sum_{t=1}^T y_t x_t - b \sum_{t=1}^T x_t^2 = 0 \end{aligned}$$

GMM and Ordinary Least Squares II

- The last line implies

$$b = \frac{\sum_{t=1}^T y_t x_t}{\sum_{t=1}^T x_t^2}$$

if $\sum_{t=1}^T x_t^2 \neq 0$

- This is just the least squares estimator
- The GMM estimator with this covariance constraint on the estimated parameter b is the same as OLS

GMM and Instrumental Variables I

- Have an equation

$$y_t = \beta x_t + \varepsilon_t$$

but

$$E[\varepsilon_t | x_t] \neq 0$$

Suppose there is another variable z_t such that

$$E[\varepsilon_t | z_t] = 0$$

- In the estimated equation

$$y_t = bx_t + e_t$$

Impose the restriction

$$E[e_t | z_t] = 0$$

GMM and Instrumental Variables II

- Again, treat conditional expectation as equivalent to linear projection, implying

$$\text{Cov} [e_t z_t] = 0$$

- Note that $\text{Cov} [e_t z_t] = 0$ imposed in the sample implies

$$\begin{aligned} & \sum_{t=1}^T e_t z_t \\ &= \sum_{t=1}^T (y_t - b x_t) z_t \\ &= \sum_{t=1}^T y_t z_t - b \sum_{t=1}^T x_t z_t = 0 \end{aligned}$$

GMM and Instrumental Variables III

- This is just the instrumental variable estimator

$$b = \frac{\sum_{t=1}^T y_t z_t}{\sum_{t=1}^T x_t z_t}$$

$$\text{if } \sum_{t=1}^T x_t z_t \neq 0$$

- This works for the identified case
- What if have two instruments for the variable x_t ?

GMM when Overidentified I

- If in

$$y_t = \beta x_t + \varepsilon_t$$

there are more restrictions than parameters and

$$E[\varepsilon_t | z_{i,t}] = 0, \quad i = 1, 2, \dots, \ell$$

cannot necessarily all be satisfied exactly

- Use a minimum distance estimator
- Suppose a model involves a set of n parameters

$$\boldsymbol{\theta}' = [\theta_1, \theta_2, \dots, \theta_n]$$

- Suppose it also involves a set of $\ell > n$ moment conditions

$$E m_i(y_t, \mathbf{x}_t, \mathbf{z}_t, \boldsymbol{\theta}) \equiv m_{i,t}(\boldsymbol{\theta}), \quad i = 1, 2, \dots, \ell$$

GMM when Overidentified II

- Then they cannot all be satisfied exactly if the equations are functionally independent
- Let the sample moment conditions be

$$\bar{m}_i = T^{-1} \sum_{t=1}^T m_{i,t}(\hat{\theta}), \quad i = 1, 2, \dots, \ell$$

- These moment conditions are set up so that the correct expected value is zero
- The moment conditions might be, for example, for $i = 1, 2$

$$m_1 = E[e_t | z_{1,t}] = 0$$

$$m_2 = E[e_t | z_{2,t}] = 0$$

GMM when Overidentified III

- One solution would be to use a minimum distance estimator to satisfy the criterion

$$q = \min_{\theta} \sum_{i=1}^{\ell} \bar{m}_i^2$$

or in matrix form

$$q = \min_{\theta} \bar{\mathbf{m}}' \bar{\mathbf{m}}$$

with

$$\bar{\mathbf{m}}' = [\bar{m}_1, \bar{m}_2, \dots, \bar{m}_{\ell}]$$

- Hansen (1982) shows that this estimator is consistent so that

$$\text{plim } \bar{m}_i = E \bar{m}_i = 0, \quad i = 1, 2, \dots, \ell$$

GMM when Overidentified IV

- He also suggests, and people invariably use, a generalization similar to weighted least squares

$$Q = \min_{\theta} \bar{\mathbf{m}}' \mathbf{W}_{\ell} \bar{\mathbf{m}}$$

where \mathbf{W}_{ℓ} is any positive-definite ℓ by ℓ matrix which can depend on the data but not on θ

- An additional assumption is that $\text{plim } \mathbf{W}_{\ell} = \mathbf{W}$
- The inverses of the individual variances on the diagonal are an obvious criterion which puts more weight on moments with less variance from the restricted value of zero
- This suggests

$$\mathbf{W} = \left[\text{Asym. Var. } \sqrt{T} \bar{\mathbf{m}} \right]^{-1}$$

which can be defined as

$$\Phi^{-1} = \mathbf{W} = \left[\text{Asym. Var. } \sqrt{T} \bar{\mathbf{m}} \right]^{-1}$$

GMM when Overidentified V

- This is the optimal weighting matrix to use
- The asymptotic variance of this GMM estimator is

$$\mathbf{V}_{GMM} = \frac{1}{T} [\mathbf{\Gamma}' \mathbf{W} \mathbf{\Gamma}]^{-1} = \frac{1}{T} [\mathbf{\Gamma}' \mathbf{\Phi}^{-1} \mathbf{\Gamma}]^{-1}$$

where $\mathbf{\Gamma}$ is the matrix of derivatives with i 'th row equal to

$$\mathbf{\Gamma}_i = \text{plim} \frac{\partial \bar{m}_i(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}'}$$

and

$$\mathbf{\Phi} = \left[\text{Asym. Var. } T^{\frac{1}{2}} \bar{m} \right]$$

- Asymptotically normally distributed

Method of Moments and Generalized Method of Moments

- Generally speaking, refer to the estimator as
 - ▶ Method of Moments when the number of moment conditions equals the number of moments
 - ▶ Generalized Method of Moments when the number of moment conditions is greater than the number of moments

Summary I

- Identification of structural equations can proceed in various ways
- Most common identification scheme for an equation is the use of exclusion restrictions
- The number of jointly endogenous variables in the equation minus one is less than or equal to the number of exogenous variables excluded from the equation
- There are a variety of estimation schemes, with Instrumental Variables (IV) and Two Stage Least Squares probably being the most common
- Full-information maximum likelihood is increasingly feasible
- GMM is very general
 - ▶ Easy to impose covariance constraints
 - ▶ IV and 2SLS are special cases, as is OLS
 - ▶ Can be one equation or many
- GMM's use
 - ▶ GMM is commonly used, especially in asset pricing, in Finance

Summary II

- ▶ GMM is commonly used when estimating first-order conditions for households and firms in Economics
- ▶ Key is orthogonality conditions
- ▶ Motivated directly by covariances
- ▶ Can be motivated by conditional linear projections
- ▶ Exposition here partly based on
 - ★ *Econometric Modelling with Time Series* by Vance Martin, Stan Hurn and David Harris
 - ★ *Econometric Analysis*, Fifth Edition by William H. Greene